

7th International Nitrogen Conference (INI 2016)

'SOLUTIONS TO IMPROVE NITROGEN USE EFFICIENCY FOR THE WORLD'



Stability of urease inhibitor added to urea

**Heitor Cantarella, Johnny R. Soares, Rafael M. Sousa,
Rafael Otto, Cleiton Sequeira**

**Agronomic Institute of Campinas (IAC), Brazil
ESALQ-USP, Piracicaba, Brazil;
Koch Agronomic Services, USA**



Common conditions for N fertilizer application in Brazil: **no till; perennials & mulch: fertilizer incorporation is difficult**



Why study the stability of NBPT

- **Urea: +60% of share of N fertilizer in Brazil.**
- **NH₃ volatilization is important: up to 20-40% losses surface-applied under high soil T and moisture**
- **Urease inhibitors sharply reduce NH₃ losses. Main inhibitor used is NBPT [N-(n-butyl thiophosphoric acid triamide)], trade name Agrotain.**
- **Shelf life of NBPT is of concern: decreased effectiveness**

OBJECTIVE: to investigate the shelf-life of NBPT-treated urea stored in different bag sizes and locations with different climates

Material and Methods





**Rondonópolis: Jan (21-32°C; x=26°C)
Jul (14-30°C; x=22°C)
Paranaguá: Jan (21-29°C; x=25°C)
Jul (14-21°C; x=17°C)**

- ✓ Urea+NBPT (870 mg NBPT/kg) stored for up to 1 year in sealed plastic bags: 500 g, 50 kg, and 750 kg (big-bags). 3 reps
- ✓ Stored in warehouses in two locations in Brazil: see map
- ✓ Sampling intervals: 1, 2, 3, 4, 5, 6, 9 and 12 months of storage
- ✓ Urea analyzed for residual NBPT and lab experiment to measure NH_3 volatilization. Red Latosol typical of Brazil (pH CaCl_2 5.5, clay: 37%)
- ✓ Controls: untreated urea and urea freshly treated with NBPT




Results



NH₃ losses and residual NBPT in urea with/without NPBP samples as a function of site and storage time (50 kg bags). P: Paranaguá; R: Rondonópolis. Bars: standard deviation.

- ❑ Residual NBPT declines with storage time
- ❑ Paranaguá (mild): NH₃ losses with UR+NBPT similar to fresh NBPT
- ❑ Rondonópolis (hot): shelf life declined after 6 months



Conclusions

- ✓ **Residual NBPT declined with storage time.**
- ✓ **Paranaguá: under similar conditions: shelf-life (SL) 9-12 months**
- ✓ **Storage at a high T (Rondonópolis) caused rapid degradation of NBPT treated urea: $SL \leq 6$ months**
- ✓ **Higher rate of NBPT in places with high T (tropical climate) may be an alternative to prolong the shelf-life**
- ✓ **Safer to restrict storage time in hot places**



See Poster 26

THANK YOU

Heitor Cantarella

cantarella@iac.sp.gov.br